

By Only

DESCRIPTION: PRENATE ELITE® is a white, oval, oil- and water-soluble, multivitamin/multimineral, film-coated tablet debossed with "PN" on one side and blank on the other. Contains no ingredient made from gluten-containing grain (wheat, barley, or rye).

Supplement Facts Serving Size: 1 Tablet			
Amount per Serving:		% Daily Value	Daily Value (Pregnant & Lactating Women)
Vitamin A (as beta-carotene)	780 mcg	90%	90%
Vitamin C (as ascorbic acid)	75 mg	125%	125%
Vitamin D ₃ (as cholecalciferol)	15 mcg	75%	75%
Vitamin E (as dl-alpha tocopheryl acetate)	18 mg	120%	120%
Thiamin (as thiamine mononitrate)	3 mg	200%	180%
Riboflavin	3.5 mg	210%	175%
Niacinamide	21 mg	105%	105%
Vitamin B ₆ (as pyridoxine HCl)	21 mg	1050%	840%
Folate (as 1.11 mg of (6S)-N5- methyltetrahydrofolic acid calcium salt (equivalent to 600 mcg of folic acid) ai	mcg DFE nd folic acid		280% mcg)
Calcium (as Formical® (calcium formate))	155 mg	20%	10%
Iron (as Sumalate® (ferrous asparto glycinat	e)) 20 mg	110%	110%
Vitamin B ₁₂ (as cyanocobalamin)	13 mcg	220%	160%
Biotin	330 mcg	110%	110%
lodine (as potassium iodide)	150 mcg	100%	100%
Magnesium (as magnesium oxide)	25 mg	10%	10%
Zinc (as zinc oxide)	15 mg	100%	100%
Copper (as cupric oxide)	1.5 mg	75%	75%

OTHER INGREDIENTS: Croscarmellose Sodium, Magnesium Stearate, Microcrystalline Cellulose, Silicon Dioxide, Stearic Acid. Film Coating (Hydroxypropyl Methylcellulose, Titanium Dioxide, Triacetin).

INDICATIONS: PRENATE ELITE® is a multivitamin/multimineral dietary supplement indicated for use in improving the nutritional status of women throughout pregnancy and in the postnatal period for both lactating and nonlactating mothers.

CONTRAINDICATIONS: PRENATE ELITE[®] is contraindicated in patients with a known hypersensitivity to any of the ingredients.

PRECAUTIONS: Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where Vitamin B₁₂ is deficient. Folic acid in doses above 1.0 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations progress.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

ADVERSE REACTIONS: Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

DOSAGE AND ADMINISTRATION: Before, during and/or after pregnancy, one tablet daily or as directed by a physician. **HOW SUPPLIED:** Bottles of 30 tablets. (75854-314-30).

The listed product number is not a National Drug Code, but has instead merely been formatted to comply with standard industry practice for pharmacy and insurance computer systems.

Store at 20° - 25°C (68° - 77°F); excursions permitted to 15° - 30°C (59° - 86°F), [See USP Controlled Room Temperature.]

MANUFACTURED FOR:

Avion Pharmaceuticals, LLC Atlanta, GA 30005 1-888-61-AVION L-0187 Rev 0620-01

THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.

Formical® is a registered trademark of Nephro-Tech 1, LLC, covered by one or more claims of U.S. Patent No. 6,528,542. Sumalate® is a registered trademark of Albion Laboratories, Inc., covered by one or more claims of U.S. Patent Nos. 5,516,925, 6,716,814, 8,007,846, and 8,425,956.